



# Exploring the Longwave Radiative Effects of Dust Aerosols

How significant is it?

*Richard A. Hansell Jr.*

*2012 Gregory G. Leptoukh On-line Giovanni Workshop*

*26 Sept 2012*

*Acknowledgements:* *S.C. Tsay, N.C. Hsu, K.N. Liou, S. Ou, J. R. Reid, Q. Ji, T.L. Roush, B.N. Holben, E.J. Welton, Z. Li, M.J. Jeong J. Pimenta, J. Haywood, O. Kalashnikova, C. Zender, M. Mishchenko, and the SMARTLabs Team*


*Hansell, R. A., et al. (2012), An assessment of the surface longwave direct radiative effect of airborne dust in Zhangye, China, during the Asian Monsoon Years field experiment (2008), J. Geophys. Res., 117, D00K39, doi:10.1029/2011JD017370*

# Outline

- ✓ *Motivation*
- ✓ *Field experiments*
- ✓ *Optical properties of dust*
- ✓ *Methodology*
- ✓ *Longwave Radiative Effects*
- ✓ *Summary*



# Motivation – Big Picture




- Aerosol radiative effects *are uncertain in climate models*
  - Dust is the most dominant aerosol by mass (Textor et al. 2006)
  - Shortwave behavior of dust is better constrained
  - Longwave effects of dust are not well known due to uncertainties in their optical properties
  - Climate models generally do not include the Longwave effects; Recent works suggest Longwave effects are important



# Here I focus on 2 field experiments to study the radiative effects of dust




- Sal Island (Cape Verde)
- $16.73^{\circ}$  N,  $22.93^{\circ}$  W
- Sea-Level
- Sept 2006

NAMMA 2006 (Zipser et al. 2006)



Mineral dust  
African Continent mixed with  
clouds during active dust period  
of the field campaign

Results published in Hansell et al. (2010)





# NASA SMARTLabs at Cape Verde (Instrumentation)




Balloon soundings



Trailers arrived on Sal Island




Aerosol & trace gases



Scanning microwave radiometer



On-site team



Active/Passive Radiation sensors


Surface-sensing **M**easurements for **A**tmospheric **R**adiative **T**ransfer  
**C**hemical, **O**ptical & **M**icrophysical **M**easurements of **I**n-Situ **T**roposphere

# AMY2008


(Lau et al. 2008;  
Li et al. 2011)

- Zhangye (desert)
- 39.0°N; 101°W
- 1.5 km ASL
- April-June 2008


Circles mark location  
of Zhangye and NASA  
SMARTlabs instruments



Ge et al. 2010





MODIS L1B RGB image 3 May 2008



Dust Parameter ( $D^*$  - Hansell et al. 2007)




# NASA SMARTLabs at Zhangye China (Instrumentation)



**CHINA<sup>2</sup>-AMY08: Cloud, Humidity Interacting Natural/  
Anthropogenic Aerosols in AMY-2008 (Asian Monsoon Years)**

# Spectral Interferometry

- Passive, fully automated, ground-based interferometer
  - Measures downward emissions
- Michelson Series MR100 (Bomem)
- 2 detectors:
  - InSb:  $3.3\mu\text{m} \leq \lambda \leq 5.5\mu\text{m}$
  - MCT:  $5.5\mu\text{m} \leq \lambda \leq 19\mu\text{m}$
- 2 Blackbody references
  - Ambient /Hot ( $60^\circ\text{C}$ )
- Scene mirror optics assembly
- Temporal frequency:  $\cong 10\text{min}$  (BB + scene + BB)
- $1\text{ cm}^{-1}$  spectral resolution





# Courtesy: NASA Giovanni System





Figure from  
Hansell et al. (2012)




Giovanni allows for rapid assessment of regional aerosol conditions

Deep Blue averaged AOT from (30 April –6 May 2008) at Zhangye during AMY.

SMARTLabs

MISR averaged AOT from (6–8 Sept 2006) at Cape Verde during NAMMA.


# Dust Physicochemical Properties: Optical Model




SEM image provided by Dr. J.S. Reid




# Dust model parameterizations



# Local measurements: model constraints




## Particle size spectra



PM10 mass concentrations








□ Optical constants of common terrestrial minerals in the thermal IR (imaginary term)

□ Large variability in absorptive peaks are evident in the single-scattering properties below


□ Dust composition plays large role in LW applications



$\alpha_{\text{ext}}$



VMD ( $\mu\text{m}$ )



□ Absorption in the IR bands

□ Spectral envelope of mass extinction efficiencies of dust versus composition, shape, and size

# Key optical parameters in thermal IR




Jeong et al. [2008]	Current study	Data source
Quartz (41%)	Quartz (43%) <sup>1</sup>	Gray (1963), Drumond (1935), Spitzer & Kleinman (1961), Philipp (1985), Longtin et al. 1988
Plagioclase (17%)	Anorthosite (17%)	J.R. Aronson and P.F.Strong [1975] - LW
K-Feldspar (9%)	Andesite (8%) <sup>1</sup>	Pollack et al. [1973] - SW
Calcite (10%)	Calcite (10%)	Long et al. [1993] - LW Marra et al. [2006] - SW
Mica (12%)	Mica (12%) <sup>2</sup>	Aronson and Strong [1975] - Muscovite (LW) Egan and Hilgeman [1979] - Illite (SW)
Chlorite (10%)	Chlorite (10%)	Mooney and Knacke [1985] - LW Thomas et al. [2009] - SW
Amphibole (1%)	Amphibole (0) <sup>1</sup>	No data available <sup>1</sup>
Dolomite (0)	Dolomite (0)	N/A <sup>3</sup>
Gypsum (0)	Gypsum (0)	N/A <sup>3</sup>
Total – 100%	Total – 100%	N/A <sup>3</sup>

Adapted from Jeong et al. 2008

Table from Hansell et al. 2012

## Zhangye

- ☐ In-situ mineralogical measurements at Zhangye
- ☐ Birefringence properties





## Cape Verde

- ☐ Transported Saharan dust at Cape Verde
- ☐ Clay , illite, kaolinite, and quartz traces
- ☐ IOR: local AERONET climatology, Volz (1973) and D' Almeida (1991)
- ☐ Birefringence properties


# Mineral composition – optical constants

# Dust Models




Zhangye

Cape Verde



# Zhangye optical model



Hematite mixture (~1%)  
SSA↓ ~ 5-10%

Larger scatter

Larger absorption



# Energy Transfer in dusty atmosphere

Top of Dust Layer

Clear-sky

Dust




Surface

Solar


Thermal Emissions

$$DARE = (I_{all-sky} \downarrow - I_{all-sky} \uparrow) - (I_{clear-sky} \downarrow - I_{clear-sky} \uparrow)$$


# Methodology



# AOT Retrievals



Zhangye



- Average visible AOT ( $0.55\mu\text{m}$ )  $\sim 0.53 \pm 0.32$ .
- Daytime/nighttime loading is comparable



# Instantaneous Surface DARE

Avg Total =  $10.1 \pm 4.9$



Avg April =  $10 \pm 7$

Avg May =  $18 \pm 8$

□ Zhangye DARE:  $\sim 2\text{-}20 \text{ Wm}^{-2}$

□ Cape Verde DARE:  $\sim 2\text{-}10 \text{ Wm}^{-2}$


Avg Total =  $6.4 \pm 2$



DARE (Zhangye)  $\sim 2\text{X}$  larger than that at Cape Verde!

□ The upper end of DARE is comparable to modeled and observed Cloud Radiative Effects ( $\geq 30 \text{ Wm}^{-2}$  - e.g., *Lockwood, 1992*); Thus it is climatically significant.


# Surface DARE (Efficiency)



- ❑ DARE (Zhangye) ranges from 31-35 Wm<sup>-2</sup>τ<sup>-1</sup>
- ❑ DARE (Taklamakan) ranges from 18-39 Wm<sup>-2</sup>τ<sup>-1</sup> (Xia et al. 2009)
- ❑ DARE (Zhangye) ~ 2X larger than that at Cape Verde

# Surface $DARE_{SW}$

- Haywood et al. 2003 estimated  $DARE_{SW} = -209 \text{ Wm}^{-2}$  during SHADE field campaign (around Cape Verde)
- Diurnally averaged  $DARE_{SW} \sim -38.4 \text{ Wm}^{-2}\tau^{-1}$  (Anderson et al. 2005)
- Considering meteorological and dust conditions to be comparable during both field studies (in September), the derived  $DARE_{LW}$  (over ocean) from NAMMA is ~42% of the diurnally averaged  $DARE_{SW}$  measured during SHADE.




$$DRE_{SW} = \frac{1}{24} \int_{\text{sunrise}}^{\text{sunset}} IDRE_{SW}(H) dH$$

- LW significance (Zhangye) ranges from 51-58% of the SW effect
- Over one half of SW cooling is compensated by LW warming
- Larger than the 33% compensation reported by Huang et al. 2009, but very close to what was found by Xia et al. (2009) - ~58%.

- *Level of significance tied to how well SSA can be constrained*



# TOA DARE (Efficiency)




DARE  $\sim 20 \text{ Wm}^{-2}\tau^{-1}$

DARE  $\sim 13 \text{ Wm}^{-2}\tau^{-1}$

- ❑ LW flux enhancement at surface is seen as a **reduction in the OLR** due to absorption by intervening dust layers
- ❑ TOA DARE  $\sim 60\%$  larger at Zhangye

# Heating Rates



- 0.25-0.30 K/day on average, with maximum heating reaching over 1.5 K/day

*Huang et al.* [2009], using CALIPSO derived vertical distributions of dust extinction over Taklamakan Desert (July 2006), reported heating rates that varied between 1-3 K/day depending on dust load, with maximum heating reaching 5.5 K/day.

# Summary Highlights

- Cape Verde: Surface  $\text{DARE}_{\text{LW}}$  varied  $\sim 2\text{-}10 \text{ Wm}^{-2}$ , with daytime/nighttime means of 6.9 and 8.4  $\text{Wm}^{-2}$ , respectively.
- Zhangye: Conservatively, surface  $\text{DARE}_{\text{LW}}$  varied about  $2\text{-}20 \text{ Wm}^{-2}$ , with daytime/nighttime means of  $\sim 12.0 \text{ Wm}^{-2}$ . Was found to be as high as  $\sim 28 \text{ Wm}^{-2}$
- Cape Verde:  $\text{DARE}_{\text{LW}}$  efficiency  $\sim 16 \text{ Wm}^{-2}\tau^{-1}$ , and nearly 42% of the diurnally averaged SW values measured during SHADE
- Zhangye:  $\text{DARE}_{\text{LW}}$  efficiency  $\sim 35 \text{ Wm}^{-2}\tau^{-1}$ , and can be as high as 58% of the diurnally averaged observed SW values.



# Summary (Cont'd)

- Cape Verde: TOA  $\text{DARE}_{\text{LW}}$  varied  $\sim 2 - 11 \text{ Wm}^{-2}$ . The  $\text{DARE}_{\text{LW}}$  efficiency at TOA is  $\sim 13 \text{ Wm}^{-2}\tau^{-1}$ .
- Zhangye: TOA  $\text{DRE}_{\text{LW}}$  varied  $\sim 2 - 16 \text{ Wm}^{-2}$ . The  $\text{DARE}_{\text{LW}}$  efficiency at TOA is  $\sim 20 \text{ Wm}^{-2}\tau^{-1}$ .
- Certainly non-negligible, the surface  $\text{DARE}_{\text{LW}}$  can be an important parameter for assessing regional changes in surface temperatures, moisture budgets, and being able to modulate the dynamics of the atmosphere.
- The upper end of DARE is comparable to Cloud Radiative Effects ( $\geq 30 \text{ Wm}^{-2}$ ); Thus it is climatically significant.
- At regional scales near dust source regions, DARE in the LW is important and can leverage the impact of SW cooling.

Thank you!